arXiv:1608.02016v1 [math.PR] 5 Aug 2016

Transporting random measures on the line and embedding excursions into Brownian motion

Günter Last*, Wenpin Tang† and Hermann Thorisson‡

August 9, 2016

Abstract

We consider two jointly stationary and ergodic random measures ξ and η on the real line $\mathbb{R}$ with equal intensities. An allocation is an equivariant random mapping from $\mathbb{R}$ to $\mathbb{R}$. We give sufficient and partially necessary conditions for the existence of allocations transporting ξ to η. An important ingredient of our approach is to introduce a transport kernel balancing ξ and η, provided these random measures are mutually singular. In the second part of the paper, we apply this result to the path decomposition of a two-sided Brownian motion into three independent pieces: a time reversed Brownian motion on $(-\infty, 0]$, an excursion distributed according to a conditional Itô's law and a Brownian motion starting after this excursion. An analogous result holds for Bismut's excursion law.

Keywords: stationary random measure, point process, allocation, invariant transport, Palm measure, shift-coupling, Brownian motion, excursion theory
AMS MSC 2010: Primary 60G57, 60G55; Secondary 60G60.

1 Introduction

In the recent papers [8, 14] and [17], balancing allocations were used for unbiased Skorohod embedding in a two-sided Brownian motion by a random time shift, and for unbiased embedding of Brownian bridge in a two-sided Brownian motion by a random space-time shift. Here 'unbiased' means that the pieces before and after the embedded pattern are independent Brownian motions. In the present paper, we extend the general theory of balancing random measures on the line so as to accommodate non-diffuse target measures. As an application, unbiased embeddings of conditional Itô's excursion laws and Bismut's excursion laws in a two-sided Brownian motion are established.

Let ξ and η be two jointly stationary and ergodic random measures on $\mathbb{R}$ with finite intensities $\lambda_\xi := \mathbb{E}\xi[0,1]$ and $\lambda_\eta := \mathbb{E}\eta[0,1]$. An *allocation* is a random (measurable) mapping $t \mapsto \tau(t)$ from $\mathbb{R}$ to $[0,\infty]$ which is equivariant under joint shifts of t and the

*Karlsruhe Institute of Technology, Germany. E-mail: guenter.last@kit.edu
†University of Berkeley, USA. E-mail: wenpintang@stat.berkeley.edu
‡University of Iceland, Iceland. E-mail: hermann@hi.is

underlying randomness; see (2.3) for an exact definition. An allocation is said to *balance* the *source* ξ and the *target* η if $\mathbb{P}(\xi(\{s \in \mathbb{R} : \tau(s) = \infty\}) > 0) = 0$ and the image measure of ξ under τ is η; that is,

$$\int \mathbf{1}\{\tau(s) \in C\}\, \xi(ds) = \eta(C), \quad C \in \mathcal{B}(\mathbb{R}),\ \mathbb{P}\text{-a.e.} \tag{1.1}$$

The balancing property (1.1) implies easily that

$$\lambda_\xi = \lambda_\eta. \tag{1.2}$$

The random variable $\tau(0)$ can be used to construct a *shift-coupling* (see [1, 19, 20]) of the Palm versions of ξ and η; see [12, 4, 9].

In this paper, we prove that if the source ξ is diffuse, and if the source and the target are mutually singular, then the equality (1.2) is not only necessary but also sufficient for the existence of a balancing allocation (note that if both ξ and η are diffuse then the $\le$ sign in (1.3) below can be replaced by $=$):

Theorem 1.1. *Assume that ξ and η are mutually singular jointly stationary and ergodic random measures on $\mathbb{R}$ such that ξ is diffuse and $\lambda_\xi = \lambda_\eta$. Then the allocation τ defined by*

$$\tau(s) := \inf\{t > s \colon \xi[s,t] \le \eta[s,t]\}, \quad s \in \mathbb{R}, \tag{1.3}$$

balances ξ and η.

In order to establish Theorem 1.1, we prove an even more general result, Theorem 3.2, which does not require ξ to be diffuse: we construct a balancing *transport kernel*, provided that $\lambda_\xi = \lambda_\eta$ and that ξ and η are mutually singular. This relies heavily on Theorem 5.1 from [8], a precursor of Theorem 1.1 where both ξ and η are assumed to be diffuse.

For further background on transports of random measures and point processes, see [11, 3, 4, 9, 8]. In [11, 3], the sources are counting and Lebesgue measures and the targets are Bernoulli and Poisson processes. In [4], the source is Lebesgue measure and the target is a simple point process, in particular a Poisson process. In [8], the source and target are both diffuse random measures, in particular local times of Brownian motion. In Theorem 1.1 above, the source is diffuse but the target is general, and according to Theorem 3.2 below (see Remark 4.2), a balancing allocation is obtained through external randomization in the case where both source and target are general. The paper [9] develops a general transport theory for random measures (on groups) with focus on transport kernels rather than only allocations.

The paper is organised as follows. Section 2 gives preliminaries on random measures, transport kernels and allocations. Section 3 provides the main transport result, Theorem 3.2. We then turn to the application to Brownian motion. Section 4 contains the key Palm and shift-coupling result for the embedding, Proposition 4.1. Section 5 is devoted to excursion theory and discusses the embedding problem. Finally, Section 6 applies Proposition 4.1 to unbiased embedding of conditional Itô's and Bismut's laws.

2 Preliminaries

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a σ-finite measure space. A *random measure* (resp. *point process*) ξ on $\mathbb{R}$ (equipped with its Borel σ-field $\mathcal{B}(\mathbb{R})$) is a kernel from Ω to $\mathbb{R}$ such that $\xi(\omega, C) < \infty$ (resp. $\xi(\omega, C) \in \mathbb{N}_0$) for $\mathbb{P}$-a.e. ω and all compact $C \subset \mathbb{R}$. We assume that $(\Omega, \mathcal{F})$ is equipped with a *measurable flow* $\theta_s \colon \Omega \to \Omega$, $s \in \mathbb{R}$. This is a family of measurable mappings such that $(\omega, s) \mapsto \theta_s\omega$ is measurable, θ_0 is the identity on Ω and

$$\theta_s \circ \theta_t = \theta_{s+t}, \quad s, t \in R, \tag{2.1}$$

where $\circ$ denotes composition.

A kernel ξ from Ω to $\mathbb{R}$ is said to be *invariant* (or *flow-adapted*) if

$$\xi(\theta_t\omega, C - t) = \xi(\omega, C), \quad C \in \mathcal{B}(\mathbb{R}), t \in \mathbb{R}, \mathbb{P}\text{-a.e. } \omega \in \Omega. \tag{2.2}$$

We assume that the measure $\mathbb{P}$ is *stationary*; that is

$$\mathbb{P} \circ \theta_s = \mathbb{P}, \quad s \in \mathbb{R},$$

where θ_s is interpreted as a mapping from $\mathcal{F}$ to $\mathcal{F}$ in the usual way:

$$\theta_s A := \{\theta_s\omega : \omega \in A\}, \quad A \in \mathcal{F},\ s \in \mathbb{R}.$$

The *invariant* σ-field $\mathcal{I} \subset \mathcal{F}$ is the class of all sets $A \in \mathcal{F}$ satisfying $\theta_s A = A$ for all $s \in \mathbb{R}$. We also assume that $\mathbb{P}$ is *ergodic*; that is for any $A \in \mathcal{I}$, we have either $\mathbb{P}(A) = 0$ or $\mathbb{P}(A^c) = 0$.

Remark 2.1. The assumption of ergodicity has been made for simplicity and can be relaxed. The assumption $\lambda_\xi = \lambda_\eta$ has then to be replaced by

$$\mathbb{E}[\xi[0,1] \mid \mathcal{I}] = \mathbb{E}[\eta[0,1] \mid \mathcal{I}], \quad \mathbb{P}\text{-a.s.}$$

We refer to [9] for more detail on this point.

A *transport kernel* is a sub-Markovian kernel K from $\Omega \times \mathbb{R}$ to $\mathbb{R}$. A transport kernel is *invariant* if

$$K(\theta_s\omega, 0, C - s) = K(\omega, s, C), \qquad s \in \mathbb{R},\ C \in \mathcal{B}(\mathbb{R}),\ \mathbb{P}\text{-a.e. } \omega \in \Omega.$$

An *allocation* [4, 9] is a measurable mapping $\tau \colon \Omega \times \mathbb{R} \to [0, \infty]$ that is *equivariant* in the sense that

$$\tau(\theta_t\omega, s - t) = \tau(\omega, s) - t, \quad s, t \in \mathbb{R},\ \mathbb{P}\text{-a.e. } \omega \in \Omega. \tag{2.3}$$

Any allocation defines a transport kernel K by $K(s, \cdot) = \mathbf{1}\{\tau(s) < \infty\}\delta_{\tau(s)}$.

Remark 2.2. In [9], a transport kernel K is Markovian; that is $K(\omega, s, \mathbb{R}) = 1$ for all $s \in \mathbb{R}$. We find it convenient to allow for $K(\omega, s, \mathbb{R}) < 1$ on an exceptional set of points (ω, s). In the same spirit, we do not assume an allocation to take on only finite values, as it is the case in [9, 8].

Let ξ and η be random measures on $\mathbb{R}$. We say that a transport kernel K *balances* ξ and η if $K(s, \mathbb{R}) = 1$ for ξ-a.e. $s \in \mathbb{R}$ and $\mathbb{P}$-a.e. and K transports ξ to η, that is,

$$\int K(s, \cdot)\, \xi(ds) = \eta, \quad \mathbb{P}\text{-a.e.} \tag{2.4}$$

If K is concentrated on an allocation τ and balances ξ and η, then we say that τ balances ξ and η.

3 Balancing mutually singular random measures

Throughout this section, let ξ and η be two invariant random measures defined on the σ-finite measure space $(\Omega, \mathcal{F}, \mathbb{P})$. We recall from the previous section that $\mathbb{P}$ is assumed to be stationary and ergodic under a given flow. In particular, the joint distribution of ξ and η is stationary and ergodic. We shall construct a transport kernel balancing ξ and η. To this end, we use the following result from [8, Theorem 5.1] in a crucial way.

Theorem 3.1. *[8] Assume that ξ and η are mutually singular diffuse invariant random measures such that $\lambda_\xi = \lambda_\eta$. Then the mapping τ defined by* (1.3) *is an allocation balancing ξ and η.*

For any $u \in [0,1]$ we define a mapping $\tau^u \colon \Omega \times \mathbb{R} \to [0,\infty]$ by

$$\tau^u(s) := \inf\{t > s \colon u\xi\{s\} + \xi(s,t) \le \eta[s,t]\}, \quad s \in \mathbb{R}, \tag{3.1}$$

where $\inf \emptyset := \infty$. Since ξ and η are invariant, it is easy to see that this is an allocation.

Theorem 3.2. *Assume that ξ and η are mutually singular invariant random measures on $\mathbb{R}$ such that $\lambda_\xi = \lambda_\eta$. Then*

$$K(s,C) := \int_0^1 \mathbf{1}\{\tau^u(s) \in C\}\, du, \quad s \in \mathbb{R},\, C \in \mathcal{B}(\mathbb{R}), \tag{3.2}$$

defines a transport kernel balancing ξ and η.

If $s \in \mathbb{R}$ satisfies $\xi\{s\} = 0$, then $\tau^u(s) = \tau^0(s)$ does not depend on $u \in [0,1]$. Therefore the kernel (3.2) reduces on $\{s : \xi\{s\} = 0\}$ to the allocation rule τ^0, that is

$$K(s,\cdot) = \mathbf{1}\{\xi\{s\} = 0\}\delta_{\tau^0(s)} + \mathbf{1}\{\xi\{s\} > 0\} \int_0^1 \mathbf{1}\{\tau^u(s) \in \cdot\}\, du. \tag{3.3}$$

If $\xi\{s\} > 0$, then we may think of $u\xi\{s\}$ as a location picked at random in the mass of ξ at s, before applying virtually the same rule τ as in Theorem 3.1. If ξ is diffuse, then $\tau^0 = \tau$, where τ is given by (1.3). Moreover, the second term on the r.h.s. of (3.3) vanishes in this case. Thus, Theorem 3.2 implies Theorem 1.1.

The proof of Theorem 3.2 relies on Theorem 3.1 and the six lemmas below. Of these lemmas all are deterministic except the final one, Lemma 3.8. For convenience we assume that ξ and η are locally finite everywhere on Ω. We shall use the decomposition $\xi = \xi^c + \xi^d$ of ξ as the sum of its diffuse part ξ^c and its purely discrete part ξ^d. The formulas $\xi^c(dt) := \mathbf{1}\{\xi\{t\} = 0\}\xi(dt)$ and $\xi^d(dt) := \mathbf{1}\{\xi\{t\} > 0\}\xi(dt)$ show that these random measures are again invariant. Similar definitions apply to η.

Theorem 3.1 assumes ξ and η to be diffuse (and mutually singular). In order to obtain a diffuse source and target, we introduce a time change. Define, for $s \in \mathbb{R}$,

$$\zeta(s) := \begin{cases} s + \xi^d[0,s) + \eta^d[0,s), & s \ge 0, \\ s - \xi^d[s,0) - \eta^d[s,0), & s < 0. \end{cases}$$

Define a random measure ξ^* on $\mathbb{R}$ by

$$\xi^*(C) := \int \mathbf{1}\{\zeta(t) \in C\}\, \xi^c(dt) + \iint \mathbf{1}\{\zeta(t) \le v \le \zeta(t) + \xi\{t\}, v \in C\}\xi\{t\}^{-1}\, dv\, \xi^d(dt).$$

Define another random measure η^* on $\mathbb{R}$ by replacing ξ with η in the above r.h.s. Then ξ^* and η^* are diffuse, and it is easy to check that these random measures are again mutually singular.

To express (ξ, η) in terms of (ξ^*, η^*), we use the generalized inverse ζ^{-1} of ζ, defined by

$$\zeta^{-1}(t) := \inf\{s \in \mathbb{R}\colon \zeta(s) \ge t\}, \quad t \in \mathbb{R}.$$

Since ζ is strictly increasing, the inverse time change ζ^{-1} is continuous.

Lemma 3.3. *Let $s \in \mathbb{R}$ and $v \in [0, \xi\{s\} \vee \eta\{s\}]$. Then $\zeta^{-1}(\zeta(s) + v) = s$.*

Proof. Since ζ is (strictly) increasing, it is easy to prove the equivalence

$$\zeta(s) \le t \iff s \le \zeta^{-1}(t), \tag{3.4}$$

valid for all $s, t \in \mathbb{R}$. Applying this to the trivial inequality $\zeta(s) \le \zeta(s) + v$ yields $s \le \zeta^{-1}(\zeta(s) + v)$. Assume by contradiction that this inequality is strict, that is, $s < s'$ where $s' = \zeta^{-1}(\zeta(s) + v)$. Trivially, $s' \le \zeta^{-1}(\zeta(s) + v)$ and (3.4) yields $\zeta(s') \le \zeta(s) + v$. This, together with $v \le \xi\{s\} \vee \eta\{s\}$, implies (for $s \ge 0$; the case $s < 0$ is similar) that

$$s' + \xi^d[0, s') + \eta^d[0, s') \le s + \xi^d[0, s] + \eta^d[0, s].$$

Now $s' > s$ and $\xi^d[0, s') + \eta^d[0, s') \ge \xi^d[0, s] + \eta^d[0, s]$, which leads to a contradiction. □

Lemma 3.4. *Let $s_1 < s_2$, $v_1 \in [0, \xi\{s_1\} \vee \eta\{s_1\}]$ and $v_2 \in [0, \xi\{s_2\} \vee \eta\{s_1\}]$. Then*

$$\xi^*[\zeta(s_1) + v_1, \zeta(s_2) + v_2] = \mathbf{1}\{\eta\{s_1\} = 0\}(\xi\{s_1\} - v_1) + \xi(s_1, s_2) + \mathbf{1}\{\eta\{s_2\} = 0\}v_2.$$

Proof. Since ξ^* is diffuse, we have

$$\xi^*[\zeta(s_1) + v_1, \zeta(s_2) + v_2] = \int \mathbf{1}\{\zeta(s_1) + v_1 < t \le \zeta(s_2) + v_2\}\, \xi^*(dt) = I_1 + I_2,$$

where

$$I_1 := \int \mathbf{1}\{\zeta(s_1) + v_1 < \zeta(s) \le \zeta(s_2) + v_2\}\, \xi^c(ds),$$

$$I_2 := \sum_{s \in \mathbb{R}} \mathbf{1}\{\xi\{s\} > 0\} \int \mathbf{1}\{\zeta(s) < v \le \zeta(s) + \xi\{s\}, \zeta(s_1) + v_1 < v \le \zeta(s_2) + v_2\}\, dv.$$

Apply first (3.4) and then Lemma 3.3 to obtain

$$I_1 = \int \mathbf{1}\{s_1 < s \le s_2\}\, \xi^c(ds) = \xi^c(s_1, s_2). \tag{3.5}$$

Turning to I_2, we restrict ourselves to the case $\eta\{s_1\} = \eta\{s_2\} = 0$. The other cases can be treated similarly. First note that the inequalities $v \le \zeta(s) + \xi\{s\}$ and $\zeta(s_1) + v_1 \le v$

imply $s_1 \le s$ (by Lemma 3.3), while the inequalities $v \le \zeta(s_2) + v_2$ and $\zeta(s) \le v$ imply $s \le s_2$. Splitting into the three cases $s = s_1,\ s_1 < s < s_2,\ s = s_2$ yields

$$\begin{aligned} I_2 =& \mathbf{1}\{\xi\{s_1\} > 0\} \int \mathbf{1}\{\zeta(s_1) + v_1 < v \le \zeta(s_1) + \xi\{s_1\}\}\, dv \\ &+ \sum_{s_1 < s < s_2} \mathbf{1}\{\xi\{s\} > 0\} \int \mathbf{1}\{\zeta(s) < v \le \zeta(s) + \xi\{s\}\}\, dv \\ &+ \mathbf{1}\{\xi\{s_2\} > 0\} \int \mathbf{1}\{\zeta(s_2) < v \le \zeta(s_2) + v_2\}\, dv. \end{aligned}$$

It follows that

$$I_2 = \xi\{s_1\} - v_1 + \xi^d(s_1, s_2) + v_2.$$

Combining this with (3.5) yields the assertion of the lemma. □

According to the following change-of-variable result, ζ^{-1} balances ξ^* and ξ.

Lemma 3.5. *Let $f\colon \mathbb{R} \to [0, \infty)$ be measurable. Then*

$$\int f(s)\, \xi(ds) = \int f(\zeta^{-1}(t))\, \xi^*(dt). \tag{3.6}$$

Proof. It suffices to establish (3.6) for $f := \mathbf{1}_{[a,b)}$, where $a < b$. Using (3.4) we obtain

$$\int \mathbf{1}_{[a,b)}(\zeta^{-1}(t))\, \xi^*(dt) = \int \mathbf{1}\{\zeta(a) \le t < \zeta(b)\}\, \xi^*(dt) = \xi[a, b),$$

where we have used Lemma 3.4 (with $v_1 = v_2 = 0$) to get the second identity. □

Define

$$\tau^*(s) := \inf\{t > s\colon \xi^*[s, t] = \eta^*[s, t]\}, \quad s \in \mathbb{R}. \tag{3.7}$$

Lemma 3.6. *Let $s \in \mathbb{R}$ with $\xi\{s\} = \eta\{s\} = 0$. Then $\tau^*(\zeta(s)) < \infty$ iff $\tau^0(s) < \infty$. In this case*

$$\zeta^{-1}(\tau^*(\zeta(s))) = \tau^0(s). \tag{3.8}$$

Proof. Take $s \in \mathbb{R}$ satisfying $\xi\{s\} = 0$. In view of the assertion (3.8) and since ξ and η are mutually singular, we can assume that $\eta\{s\} = 0$. We abbreviate $t^* := \tau^*(\zeta(s))$.

First consider the case $t^* = \zeta(s)$. Then $\zeta^{-1}(t^*) = s$ (by Lemma 3.3) and we need to show that $\tau^0(s) = s$. There are $t_n > t^*$, $n \in \mathbb{N}$, such that $\xi^*[\zeta(s), t_n] = \eta^*[\zeta(s), t_n]$ and $t_n \downarrow t^*$. We distinguish two cases. In the first case, there are infinitely many $n \in \mathbb{N}$ such that $t_n = \zeta(s_n) + v_n$ for some $s_n \in \mathbb{R}$ satisfying $\eta\{s_n\} = 0$ and $v_n \in [0, \xi\{s_n\}]$. Lemma 3.4 implies $\xi[s, s_n) + v_n = \eta[s, s_n]$ and hence $\xi[s, s_n) \le \eta[s, s_n]$. Since $t^* < t_n = \zeta(s_n + v_n)$ we obtain from (3.4) and Lemma 3.3 that $s_n > \zeta^{-1}(t^*) = s$. Lemma 3.3 and the continuity of ζ^{-1} imply $s_n = \zeta^{-1}(t_n) \downarrow \zeta^{-1}(t^*) = s$ along the chosen subsequence. Hence $\tau^0(s) = s$. In the second case, there are infinitely many $n \in \mathbb{N}$ such that $t_n = \zeta(s_n) + v_n$ for some $s_n \in \mathbb{R}$ satisfying $\eta\{s_n\} > 0$ and $v_n \in [0, \eta\{s_n\}]$. Then $\xi\{s_n\} = 0$ and Lemma 3.4 implies

$\xi[s, s_n) = \eta[s, s_n) + v_n$ and hence $\xi[s, s_n) \le \eta[s, s_n]$. As before, it follows that $s_n > s$ and $s_n \downarrow s$ along the chosen subsequence. Hence $\tau^0(s) = s$ in this case.

In the remainder of the proof, we assume that $t^* > \zeta(s)$. By definition,

$$\xi^*[\zeta(s), r] > \eta^*[\zeta(s), r], \quad \zeta(s) < r < t^*, \tag{3.9}$$

as well as

$$\xi^*[\zeta(s), t^*] = \eta^*[\zeta(s), t^*]. \tag{3.10}$$

Assume first that $t^* = \zeta(t) + v$ for some $t \ge s$ with $\eta\{t\} = 0$ and $v \in [0, \xi\{t\}]$. Then $t > s$ (by (3.10)) and Lemma 3.3 implies that $\zeta^{-1}(t^*) = t$. We want to show that $\tau^0(s) = t$. Let $t' \in (s, t)$. If $\xi\{t'\} = 0$, we set $r := \zeta(t') + \eta\{t'\}$. Then $r < \zeta(t) \le t^*$ and (3.9) together with Lemma 3.4 imply that $\xi[s, t') > \eta[s, t']$. If $\xi\{t'\} > 0$, we set $r := \zeta(t')$ to obtain the same inequality and hence

$$\xi[s, t') > \eta[s, t'], \quad s < t' < t. \tag{3.11}$$

On the other hand, we have from (3.10) and Lemma 3.4 that $\xi[s, t) + v = \eta[s, t]$, so that $\xi[s, t) \le \eta[s, t]$. Hence $\tau^0(s) = t = \zeta^{-1}(t^*)$ and (3.8) follows.

The second possible case is $t^* = \zeta(t) + v$ for some $t \ge s$ with $\eta\{t\} > 0$ and $v \in [0, \eta\{t\}]$. Since ξ and η are mutually singular, we have $\xi\{t\} = 0$. Again this implies $t > s$ and (3.11). Lemma 3.4 implies that $\xi[s, t) + v = \eta[s, t)$ and hence $\xi[s, t) \le \eta[s, t]$. Therefore $\tau^0(s) = t = \zeta^{-1}(t^*)$, where we have used Lemma 3.3.

Assume, finally, that $t^* = \infty$, so that (3.9) holds for all $r > \zeta(s)$. Let $t' > s$. If $\xi\{t'\} = \eta\{t'\} = 0$, we take $r = \zeta(t')$ to obtain from Lemma 3.4 that $\xi[s, t') > \eta^*[\zeta(s), t']$. If $\xi\{t'\} > 0$ (and hence $\eta\{t'\} = 0$), we take $r = \zeta(t')$ to obtain from Lemma 3.4 that $\xi[s, t') > \eta[s, t']$. If $\eta\{t'\} > 0$ (and hence $\xi\{t'\} = 0$), we take $r = \zeta(t') + \eta\{t'\}$ to obtain from Lemma 3.4 that $\xi[s, t') > \eta[s, t']$. Hence $\tau^0(s) = \infty$. □

Lemma 3.7. *Let $s \in \mathbb{R}$ with $\xi\{s\} > 0$ and $u \in [0, 1)$. Then $\tau^*(\zeta(s) + u\xi\{s\}) < \infty$ iff $\tau^{1-u}(s) < \infty$. In this case*

$$\zeta^{-1}(\tau^*(\zeta(s) + u\xi\{s\})) = \tau^{1-u}(s). \tag{3.12}$$

Proof. Since ξ and η are mutually singular we have $\eta\{s\} = 0$. Moreover, Since $u < 1$ we have $\eta[s, s + \varepsilon] < (1 - u)\xi\{s\}$ for all sufficiently small $\varepsilon > 0$ and therefore $\tau^{1-u}(s) > s$ and (by Lemma 3.4) $\tau^*(\zeta(s) + u\xi\{s\}) > \zeta(s)$. The proof can now proceed similar to that of Lemma 3.6. The main tool is again Lemma 3.4. In contrast to the case $\xi\{s\} = 0$, it has to be applied with $s_1 = s$ and $v_1 = u\xi\{s\}$. Further details are omitted. □

In the upcoming proof of Theorem 3.2, we will use that τ^* balances ξ^* and η^*. Since ξ^* and η^* need not be jointly stationary, Theorem 3.1 cannot be used directly to establish this fact. As an intermediate step, we need the following lemma which presents (shifted and length-biased) versions of ξ^* and η^* that Theorem 3.1 can be applied to. We abuse notation by introducing for any measure μ on $\mathbb{R}$ and $s \in \mathbb{R}$ a new measure $\theta_s\mu$ by $\theta_s\mu := \mu(\cdot + s)$. If μ' is another measure on $\mathbb{R}$, we write $\theta_s(\mu, \mu') := (\theta_s\mu, \theta_s\mu')$.

Lemma 3.8. *Let ξ and η be random measures on $\mathbb{R}$ defined on $(\Omega, \mathcal{F}, \mathbb{P})$ and let ξ^* and η^* be as above. Extend $(\Omega, \mathcal{F}, \mathbb{P})$ to support a random variable U that is uniform on $[0,1)$ conditionally on (ξ^*, η^*). Let $S_1 = 1 + \xi^d[0,1) + \eta^d[0,1)$ and define a σ-finite measure $\mathbb{P}^*$ on $(\Omega, \mathcal{F})$ by $d\mathbb{P}^* = S_1 d\mathbb{P}$. Then $\theta_{US_1}(\xi^*, \eta^*)$ is stationary and ergodic under $\mathbb{P}^*$ and the intensities are*

$$\mathbb{E}^*(\theta_{US_1}\xi^*)[0,1) = \mathbb{E}\xi[0,1), \quad \mathbb{E}^*(\theta_{US_1}\eta^*)[0,1) = \mathbb{E}\eta[0,1).$$

Proof. Let M denote the space of locally finite measures on $\mathbb{R}$, equipped with the natural Kolmogorov σ-field. We have assumed that ξ and η are random elements of M.

Take $t \geq 0$ and let $f\colon M \times M \to [0,\infty)$ be bounded and measurable. For $n \in \mathbb{N}$, let $A_n \in \mathcal{F}$ be such that $\mathbb{P}(A_n) < \infty$ and $A_1 \subset A_2 \subset A_3 \subset \ldots \uparrow \Omega$. Set $f_n = f1_{A_n}$ and note that $\mathbb{P}(A_n) < \infty$ implies that $\mathbb{E}\int_{S_1}^{S_1+t} f_n(\theta_s(\xi^*, \eta^*))ds$ is finite. Since $\int_t^{S_1} f_n(\theta_s(\xi^*, \eta^*))ds$ could be negative, the finiteness is needed for the last equality in

$$\begin{aligned}\mathbb{E}^* f_n(\theta_t\theta_{US_1}(\xi^*, \eta^*)) &= \mathbb{E}S_1 f_n(\theta_t\theta_{US_1}(\xi^*, \eta^*)) = \mathbb{E}\int_t^{S_1+t} f_n(\theta_s(\xi^*, \eta^*))ds \\ &= \mathbb{E}\int_t^{S_1} f_n(\theta_s(\xi^*, \eta^*))ds + \mathbb{E}\int_{S_1}^{S_1+t} f_n(\theta_s(\xi^*, \eta^*))ds.\end{aligned}$$

Now note that $\theta_{S_1}(\xi^*, \eta^*)$ is the same measurable mapping of $\theta_1(\xi, \eta)$ as (ξ^*, η^*) is of (ξ, η). Since $\theta_1(\xi, \eta)$ has the same $\mathbb{P}$-distribution as (ξ, η), this implies that $\theta_{S_1}(\xi^*, \eta^*)$ has the same $\mathbb{P}$-distribution as (ξ^*, η^*). This further yields the first equality in

$$\begin{aligned}\mathbb{E}^* f_n(\theta_t\theta_{US_1}(\xi^*, \eta^*)) &= \mathbb{E}\int_t^{S_1} f_n(\theta_s(\xi^*, \eta^*))ds + \mathbb{E}\int_0^t f_n(\theta_s(\xi^*, \eta^*))ds \\ &= \mathbb{E}\int_0^{S_1} f_n(\theta_s(\xi^*, \eta^*))ds.\end{aligned}$$

Send $n \to \infty$ and use the monotone convergence theorem to obtain

$$\mathbb{E}^* f(\theta_t\theta_{US_1}(\xi^*, \eta^*)) = \mathbb{E}\int_0^{S_1} f(\theta_s(\xi^*, \eta^*))ds.$$

So $\mathbb{E}^* f(\theta_t\theta_{US_1}(\xi^*, \eta^*))$ does not depend on t; that is, $\theta_{US_1}(\xi^*, \eta^*)$ is stationary under $\mathbb{P}^*$.

Next we prove the ergodicity assertion. It is then not hard to prove that

$$((\theta_t\xi)^*, (\theta_t\eta)^*) = (\theta_{\zeta(t)}\xi^*, \theta_{\zeta(t)}\eta^*), \quad t \in \mathbb{R}. \tag{3.13}$$

Let $A \subset M \times M$ be a measurable set that is invariant under diagonal shifts. Then (3.13) shows for all $(\omega, t) \in \Omega \times \mathbb{R}$ that $(\xi(\theta_t\omega)^*, \eta(\theta_t\omega)^*) \in A$ if and only if $(\xi(\omega)^*, \eta(\omega)^*) \in A$. Since $\mathbb{P}$ is ergodic, we obtain that either $\mathbb{P}((\xi^*, \eta^*) \in A) = 0$ or $\mathbb{P}((\xi^*, \eta^*) \notin A) = 0$. Therefore, the ergodicity of the shifted length-biased version of (ξ^*, η^*) follows from the two facts that the zero sets of $\mathbb{P}^*$ are the same as those of $\mathbb{P}$, and that randomly shifted invariant sets remain invariant.

It remains to prove the intensity result. Let λ be Lebesgue measure on $\mathbb{R}$. Since λ is shift-invariant and $\theta_{S_1}\xi^*$ has the same $\mathbb{P}$-distribution as ξ^*, we have

$$\mathbb{E}(\lambda \otimes \xi^*)(A) = \mathbb{E}(\lambda \otimes \xi^*)((S_1, S_1) + A), \quad A \in \mathcal{B}(\mathbb{R}^2).$$



Figure 1: The region $\{(s,x) : 0 \le s < S_1, s \le x < s+1\}$ has the same Lebesgue measure as $(\{(s,x) : 0 \le x < S_1, x-1 < s \le x\} \setminus A) \cup ((S_1, S_1) + A)$.

Apply this with $A = \{(s,x) : -1 \le s < 0, 0 \le x < s+1\}$ to obtain

$$\begin{aligned}
\mathbb{E}^*(\theta_{US_1}\xi^*)[0,1) &= \mathbb{E}S_1 \int_0^1 (\theta_{uS_1}\xi^*)[0,1)du = \mathbb{E}\int_0^{S_1} \xi^*(s+[0,1))ds \\
&= \mathbb{E}(\lambda \otimes \xi^*)(\{(s,x) : 0 \le s < S_1, s \le x < s+1\}) \\
&= \mathbb{E}(\lambda \otimes \xi^*)((\{(s,x) : 0 \le x < S_1, x-1 < s \le x\} \setminus A) \cup ((S_1, S_1) + A)) \\
&= \mathbb{E}(\lambda \otimes \xi^*)(\{(s,x) : 0 \le x < S_1, x-1 < s \le x\}) \\
&\qquad - \mathbb{E}(\lambda \otimes \xi^*)(A) + \mathbb{E}(\lambda \otimes \xi^*)((S_1, S_1) + A)) \\
&= \mathbb{E}(\lambda \otimes \xi^*)(\{(s,x) : 0 \le x < S_1, x-1 < s < x\}) \\
&= \mathbb{E}\xi^*[0, S_1) = \mathbb{E}\xi[0,1).
\end{aligned}$$

In the same way, we obtain $\mathbb{E}^*(\theta_{US_1}\eta^*)[0,1) = \mathbb{E}\eta[0,1)$. □

Proof of Theorem 3.2. By Theorem 3.1 and Lemma 3.8, the random mapping $\tau' \colon \mathbb{R} \to [0,\infty]$ defined by

$$\tau'(s) := \inf\{t > s \colon (\theta_{US_1}\xi^*)[s,t] \le (\theta_{US_1}\eta^*)[s,t]\}, \quad s \in \mathbb{R},$$

balances $\theta_{US_1}\xi^*$ and $\theta_{US_1}\eta^*$ under $\mathbb{P}^*$. Since $\mathbb{P}^*$ and $\mathbb{P}$ are equivalent, this implies that τ' balances $\theta_{US_1}\xi^*$ and $\theta_{US_1}\eta^*$ under $\mathbb{P}$. This remains true if the origin is shifted to some random location. Recall the definition (3.7) of τ^*. Shift the origin to $-US_1$ to obtain

$$\tau^* = \tau'(\cdot - US_1) + US_1, \qquad \xi^* = \theta_{-US_1}\theta_{US_1}\xi^*, \qquad \eta^* = \theta_{-US_1}\theta_{US_1}\eta^*.$$

Thus, τ^* balances ξ^* and η^* under $\mathbb{P}$. We assume for simplicity that

$$\int \mathbf{1}\{\tau^*(s) \in \cdot\}\, \xi^*(ds) = \eta^* \tag{3.14}$$

holds everywhere on Ω.

We want to prove that

$$\iint f(t)\,K(s,dt)\,\xi(ds) = \int f(s)\,\eta(ds) \tag{3.15}$$

for all measurable $f\colon \mathbb{R} \to [0,\infty)$, implying the theorem. Applying Lemma 3.5 with η in place of ξ and then the balancing property (3.14) of τ^* yields

$$\int f(t)\,\eta(dt) = \int f(\zeta^{-1}(t))\,\eta^*(dt) = \int \mathbf{1}\{\tau^*(s) < \infty\} f(\zeta^{-1}(\tau^*(s)))\,\xi^*(ds) = I_1 + I_2,$$

where

$$I_1 := \int \mathbf{1}\{\tau^*(\zeta(s)) < \infty\} f(\zeta^{-1}(\tau^*(\zeta(s)))\,\xi^c(ds),$$

$$I_2 := \iint \mathbf{1}\{0 \le u \le 1\}\mathbf{1}\{\tau^*(\zeta(s) + u\xi\{s\})) < \infty\} f(\zeta^{-1}(\tau^*(\zeta(s) + u\xi\{s\})\,du\,\xi^d(ds).$$

Here we have used the definition of ξ^* and a change of variables in the second summand.

Lemma 3.6 implies that $I_1 = \int \mathbf{1}\{\tau^0(s) < \infty\} f(\tau^0(s))\,\xi^c(ds)$. Note that $\tau^u(s) = \tau^0(s)$ for all $u \in [0,1]$, whenever $\xi\{s\} = 0$. It follows that

$$I_1 = \iint f(t)K(s,dt)\,\xi^c(ds).$$

To treat I_2, we use Lemma 3.7 to obtain that

$$I_2 = \iint \mathbf{1}\{0 \le u < 1\}\mathbf{1}\{\tau^{1-u}(s) < \infty\} f(\tau^{1-u}(s))\,du\,\xi^d(ds) = \iint f(t)K(s,dt)\,\xi^d(ds),$$

which proves (3.15). □

4 On Palm measures and shift-coupling

Let ξ be an invariant random measure on $\mathbb{R}$. The *Palm measure* $\mathbb{P}_\xi$ of ξ (with respect to $\mathbb{P}$) is defined by

$$\mathbb{P}_\xi(A) := \mathbb{E}_\mathbb{P} \int \mathbf{1}_{[0,1]}(s)\mathbf{1}_A(\theta_s)\,\xi(ds), \quad A \in \mathcal{F}. \tag{4.1}$$

This is a σ-finite measure on $(\Omega, \mathcal{F})$ satisfying the *refined Campbell formula*

$$\iint f(\theta_s\omega, s)\,\xi(\omega, ds)\,\mathbb{P}(d\omega) = \iint f(\omega, s)\,ds\,\mathbb{P}_\xi(d\omega) \tag{4.2}$$

for each measurable $f\colon \Omega \times \mathbb{R} \to [0,\infty)$, where $\mathbb{E}$ refers to the integration with respect to $\mathbb{P}$. If the *intensity* $\mathbb{P}_\xi(\Omega)$ of ξ is positive and finite, then $\mathbb{P}_\xi$ can be normalized to yield the Palm probability measure of ξ. This normalization can be interpreted as conditional version of $\mathbb{P}$ given that the origin represents a point randomly chosen in the mass of ξ; see [9, 10].

The following *shift-coupling* result is a consequence of Theorem 3.2 and [9, Theorem 4.1].

Proposition 4.1. *Let ξ and η satisfy the assumptions of Theorem 3.2 and define the allocation maps τ^u, $u \in [0,1]$, by (3.1). Then*

$$\int_0^1 \mathbb{P}_\xi(\theta_{\tau^u} \in \cdot)\, du = \mathbb{P}_\eta, \tag{4.3}$$

where $\theta_{\tau^u} \colon \Omega \to \Omega$ is defined by $\theta_{\tau^u}(\omega) := \theta_{\tau^u(\omega,0)}\omega$, $\omega \in \Omega$. If, moreover, ξ is diffuse then

$$\mathbb{P}_\xi(\theta_\tau \in \cdot) = \mathbb{P}_\eta, \tag{4.4}$$

where the allocation map τ is defined by (1.3).

Remark 4.2. If we extend $(\Omega, \mathcal{F}, \mathbb{P})$ to support an independent random variable U uniformly distributed on $[0,1]$, then τ^U is a *randomized* allocation balancing ξ and η even when ξ is not diffuse. Moreover, (4.3) can then be written in the same shift-coupling form as (4.4), namely

$$\mathbb{P}_\xi(\theta_{\tau^U} \in \cdot) = \mathbb{P}_\eta.$$

In Brownian excursion theory, it is natural to define Palm measures of random measures that are not locally finite. A *σ-finite random measure* ξ is a kernel from Ω to $\mathbb{R}$ with the following property. There exist measurable sets $A_n \in \mathcal{F}$, $n \in \mathbb{N}$, such that $\cup_n A_n = \Omega$ and

$$\int \mathbf{1}\{s \in \cdot, \theta_s \in A_n\}\, \xi(ds) \tag{4.5}$$

is a random measure for each $n \in \mathbb{N}$. In this case, the Palm measure $\mathbb{P}_\xi$ can again be defined by (4.1). It is σ-finite and satisfies the refined Campbell formula (4.2). We shall use such a measure in Proposition 6.1.

5 Excursions of Brownian motion

In the next two sections, we assume that Ω is the class of all continuous functions $\omega \colon \mathbb{R} \to \mathbb{R}$ equipped with the Kolmogorov product σ-algebra $\mathcal{F}$. Let $B = (B_t)_{t\in\mathbb{R}}$ denote the identity on Ω. The flow is given by

$$(\theta_t\omega)_s := \omega_{t+s}. \tag{5.1}$$

Let $\mathbb{P}_0$ denote the distribution of a two-sided standard Brownian motion. Define $\mathbb{P}_x := \mathbb{P}_0(B + x \in \cdot)$, $x \in \mathbb{R}$, and the σ-finite and stationary measure

$$\mathbb{P} := \int \mathbb{P}_x\, dx. \tag{5.2}$$

By [8, Theorem 3.5] this $\mathbb{P}$ is ergodic. Expectations with respect to $\mathbb{P}_x$ and $\mathbb{P}$ are denoted by $\mathbb{E}_x$ and $\mathbb{E}_\mathbb{P}$, respectively.

Let $t \in \mathbb{R}$. For each real-valued function w whose domain contains $[t, \infty)$, let

$$D_t(w) := \inf\{s > t : w(s) = 0\},$$

where $\inf \emptyset := \infty$. We abbreviate $D(w) := D_0(w)$. Then

$$R_t := D(\theta_t B) = D_t(B) - t$$

is the time taken by B to hit 0 (starting at time t), while

$$L := \{t \in \mathbb{R} : R_{t-} = 0, R_t > 0\},$$

is the set of left ends of *excursion intervals*. The space E of excursions is the class of all continuous functions $e\colon [0,\infty) \to \mathbb{R}$ such that $e(0) = 0$, $0 < D(e) < \infty$, and $e(t) = 0$ for all $t \geq D(e)$. The number $D(e)$ is called the lifetime of the excursion. We equip E with the Kolmogorov product σ-field $\mathcal{E}$. For $s \in L$, define the (random) excursion $\epsilon_s \in E$ starting at time s by

$$\epsilon_s(t) := \begin{cases} B_{s+t}, & \text{if } 0 \leq t \leq R_s, \\ 0, & \text{if } t > R_s. \end{cases}$$

It is convenient to introduce the function $\delta \equiv 0$ on $[0,\infty)$, and to define $\epsilon_s := \delta$ for $s \notin L$. Then $(\omega, t) \mapsto \epsilon_t(\omega)$ is a measurable mapping with values in $E_\delta := E \cup \{\delta\}$. Note that $D(\delta) = 0$.

Define a σ-finite invariant random measure N on $\mathbb{R}$ by

$$N(C) := \sum_{s \in L} \mathbf{1}\{s \in C\}, \quad C \in \mathcal{B}(\mathbb{R}). \tag{5.3}$$

Here the invariance is obvious, while we may choose $A_1 := \{D \in \{0,\infty\}\}$ and $A_n := \{D \geq 1/n\}$, $n \geq 2$, in (4.5), to see that N is σ-finite. It follows from the refined Campbell formula (4.2) that

$$\mathbb{E} \int \mathbf{1}\{(s, \epsilon_s) \in \cdot\}\, N(ds) = \iint \mathbf{1}\{(s,e) \in \cdot\}\, ds\, \nu(de),$$

where $\nu := \mathbb{P}_N(\epsilon_0 \in \cdot)$, that is

$$\nu(A) = \mathbb{E} \int_{[0,1]} \mathbf{1}\{\epsilon_s \in A\}\, N(ds), \quad A \in \mathcal{E}. \tag{5.4}$$

(Note that $\nu\{\delta\} = 0$.) In fact, Pitman [16] showed that ν coincides with *Itô's excursion law* (suitably normalized).

In the next section, we are interested in finding random patterns distributed according to a conditional Itô's excursion law in an unbiased way. Before embarking on this by means of Palm and transport theory, we check what happens if we simply choose for a given $A \in \mathcal{E}$ the first A-excursion (an excursion belonging to A) to the right of the origin. For a measure μ and a set A such that $0 < \mu(A) < \infty$, we define the conditional measure $\mu(\cdot \mid A) = \mu(\cdot \cap A)/\mu(A)$. The following well-known result can be derived with the help of excursion theory; see [18, Lemma XII(1.13)]. An alternative proof can be based on Neveu's exchange formula and the regenerative structure described below.

Proposition 5.1. *Let $A \in \mathcal{E}$ satisfy $0 < \nu(A) < \infty$ and define*

$$S_A := \inf\{t > 0 : t \in L, \epsilon_t \in A\}.$$

Then $\mathbb{P}_0(\epsilon_{S_A} \in \cdot) = \nu(\cdot \mid A)$.

Thus, the first A-excursion to the right of the origin of a standard Brownian motion B is distributed according to the conditional Itô's law. Also, with L_A the length of the A-excursion, an independent standard Brownian motion $(\theta_{S_A+L_A}B)_{t\geq 0}$ starts at its right endpoint. However, the process $(\theta_{S_A}B)_{t\leq 0}$ is not a standard Brownian motion because it starts with a path segment of positive length S_A without an A-excursion and then at time S_A an independent standard Brownian motion starts. In fact, it follows from excursion theory (in particular from the Poisson nature of the point process (6.6)) that both $\ell^0([0, S_A])$ and $\ell^0([S'_A, 0])$ have (under $\mathbb{P}_0$) an exponential distribution with rate parameter $\nu(A)$, where $S'_A := \sup\{t < 0 : t \in L, \epsilon_t \in A\}$. Since these random variables are independent, it follows that $S'_A(\theta_{S_A}B)$ has a Gamma distribution with shape parameter 2. Hence the embedding of the Itô law in Proposition 5.1 is not unbiased.



Figure 2: The times S_A and $S'_A(\theta_{S_A}B)$ in a two-sided Brownian motion.

The above description suggests that a two-sided standard Brownian motion has the following structure. It splits into a two-sided sequence of independent segments such that: (1) every odd-numbered segment is an A-excursion distributed according to the conditional Itô's law; (2) every even-numbered segment except the one enumerated 0 is a standard Brownian motion starting from zero running until the first time that an A-excursion occurs; (3) the segment enumerated 0 consists of two independent segments of that type. In addition to this, the origin of the standard Brownian motion should be placed at random in the segment enumerated 0 according to the local time at zero.

This structure finally suggests that since the Palm version with respect to the point process N_A is point-stationary, it should have the above structure except that the segment enumerated 0 should be like the other even-numbered segments and should end at time 0. In particular, this would imply that under $\mathbb{P}_{N_A}$, the backward Brownian motion $(B_t)_{t\leq 0}$ and the excursion ϵ_0 beginning at 0 are independent.

6 Finding an extra excursion

Let A be a measurable set of excursions with positive and finite Itô measure. In this section, we will use Proposition 4.1 (with $\xi = \ell^0$) for unbiased embedding of an A-excursion; that is, in the terminology of [3] and [4], for finding an *extra A-excursion* in the sense that if we shift the origin to that excursion and then remove it, then the process remains a standard Brownian motion. For this purpose, we need to show that under the Palm measure of N, the Brownian motion is decomposed into three independent pieces: a time reversed Brownian motion on $(-\infty, 0]$, an excursion with 'distribution' ν and a Brownian motion starting after this excursion. A formal statement of this result requires some notation.

Let Ω^- (resp. Ω^+) denote the space of all continuous functions w on $(-\infty, 0]$ (resp. $[0, \infty)$) with $w(0) = 0$. The *concatenation* of $w_1 \in \Omega^-$,$w_2 \in E$ and $w_3 \in \Omega^+$ is the function $w_1 \odot w_2 \odot w_3 \in \Omega$ defined by

$$w_1 \odot w_2 \odot w_3(t) := \begin{cases} w_1(t), & \text{if } t \le 0, \\ w_2(t), & \text{if } 0 < t < D(w_2), \\ w_3(t - D(w_2)), & \text{if } D(w_2) \le t. \end{cases}$$

The *concatenation* of σ-finite measure ν_1 on Ω^-, ν_2 on E, and ν_3 on Ω^+ is the measure $\nu_1 \odot \nu_2 \odot \nu_3$ on Ω defined by

$$\nu_1 \odot \nu_2 \odot \nu_3 := \iiint \mathbf{1}\{w_1 \odot w_2 \odot w_3 \in \cdot\}\, \nu_1(dw_1)\, \nu_2(dw_2)\, \nu_3(dw_3).$$

Let $\mathbb{P}^-$ (resp. $\mathbb{P}^+$) denote the law of $B^- := (B_t)_{t \le 0}$ (resp. $B^+ := (B_t)_{t \ge 0}$).

The following proposition extends a result from Pitman [16].

Proposition 6.1. *The Palm measure of N is given by $\mathbb{P}_N = \mathbb{P}^- \odot \nu \odot \mathbb{P}^+$.*

Let $A \in \mathcal{E}$ be such that $0 < \nu(A) < \infty$. Define an invariant random measure N_A on $\mathbb{R}$ by

$$N_A(C) := \frac{1}{\nu(A)} \int \mathbf{1}\{s \in C, \epsilon_s \in A\}\, N(ds), \quad C \in \mathcal{B}(\mathbb{R}), \tag{6.1}$$

We then have the following immediate consequence of Proposition 6.1.

Corollary 6.2. *Let $A \in \mathcal{E}$ satisfy $0 < \nu(A) < \infty$ and define the point process N_A by (6.1). Then the Palm measure of N_A is given by $\mathbb{P}_N = \mathbb{P}^- \odot \nu(\cdot|A) \odot \mathbb{P}^+$.*

In order to prove Proposition 6.1 and then to apply Proposition 4.1, we use the local times of B. For $x \in \mathbb{R}$, we denote by ℓ^x the random measure associated with the *local time* of B at $x \in \mathbb{R}$ (under $\mathbb{P}_0$). The global construction in [15] (see also [6, Proposition 22.12] and [13, Theorem 6.43]) guarantees the existence of a version of local times with the following properties. The random measure ℓ^0 is $\mathbb{P}_x$-a.e. diffuse for any $x \in \mathbb{R}$ and

$$\ell^0(\theta_t\omega, C - t) = \ell^0(\omega, C), \quad C \in \mathcal{B}(\mathbb{R}),\, t \in \mathbb{R},\, \mathbb{P}_x\text{-a.s.},\, x \in \mathbb{R}, \tag{6.2}$$

$$\ell^y(\omega, \cdot) = \ell^0(\omega - y, \cdot), \quad \omega \in \Omega,\, y \in \mathbb{R}, \tag{6.3}$$

$$\operatorname{supp} \ell^x(\omega) = \{t \in \mathbb{R} : B_t = x\}, \quad \omega \in \Omega,\, x \in \mathbb{R}, \tag{6.4}$$

where $\operatorname{supp}\mu$ is the support of a measure μ on $\mathbb{R}$. Equation (6.3) implies that ℓ^y is $\mathbb{P}_x$-a.e. diffuse for any $x \in \mathbb{R}$ and is invariant in the sense of (6.2). By a classical result from [2] (see also [8, Lemma 2.3]), the Palm measure of ℓ^x is given by

$$\mathbb{P}_{\ell^x} = \mathbb{P}_x, \quad x \in \mathbb{R}. \tag{6.5}$$

The following path decomposition of a two-sided Brownian motion is the main result of this section.

Theorem 6.3. *Let $A \in \mathcal{E}$ be such that $0 < \nu(A) < \infty$ and define the random measure N_A by (6.1). Let*

$$T := \inf\{t > 0 : \ell^0[0,t] \le N_A[0,t]\}.$$

Then $\mathbb{P}_0(T < \infty) = 1$ and $\mathbb{P}_0(\theta_T B \in \cdot) = \mathbb{P}^- \odot \nu(\cdot|A) \odot \mathbb{P}^+$.

Proof of Proposition 6.1. For $t \ge 0$, let $\ell^0_t := \ell^0([0,t])$. Define the right inverse of ℓ^0 by $\tau_s := \inf\{t \ge 0 : \ell^0_t > s\}$ for $s \ge 0$. By (6.4),

$$\{t \ge 0 : B_t \ne 0\} = \bigcup_{s>0} (\tau_{s-}, \tau_s).$$

If $\tau_{s-} < \tau_s$, then (τ_{s-}, τ_s) is an excursion interval away from 0. A classical result of Itô [5] (see also [6, Theorem 22.11] and [18, Theorem XII(2.4)]) shows that the random measure

$$\Phi := \sum_{s>0:\tau_{s-}<\tau_s} \delta_{(s,\epsilon_{\tau_{s-}})} \tag{6.6}$$

is a Poisson process on $(0,\infty) \times E$ under $\mathbb{P}_0$ with intensity measure

$$\mathbb{E}_0 \sum_{s>0:\tau_{s-}<\tau_s} \mathbf{1}\{(s,\epsilon_{\tau_{s-}}) \in \cdot\} = \int_E \int_0^\infty \mathbf{1}\{(s,e) \in \cdot\}\, ds\, \nu(de).$$

The excursion law ν satisfies

$$\nu(D \in dr) = (2\pi)^{-1/2} r^{-3/2}\, dr; \tag{6.7}$$

see [18, Section XII.2] or [6, Theorem 22.5].

For the purpose of this proof, it is convenient to enlarge for each $x \in \mathbb{R}$ the probability space $(\Omega, \mathcal{F}, \mathbb{P}_x)$ to a probability space $(\Omega', \mathcal{F}', \mathbb{P}'_x)$, so as to support a Poisson process Φ' on $(0,\infty) \times E$ with intensity measure $dt\nu(de)$, independent of B. We say that an event occurs $\mathbb{P}'$-a.e., if it occurs $\mathbb{P}'_x$-a.e. for every $x \in \mathbb{R}$. Define,

$$\tau'(t) := \int \mathbf{1}\{s \le t\} D(e)\, \Phi'(d(s,e)), \quad t \ge 0. \tag{6.8}$$

By (6.7), $\int \min\{D(e), 1\}\, \nu(de) < \infty$. Hence [6, Lemma 12.13] (see also [7, Proposition 12.1]) shows that (6.8) converges $\mathbb{P}'$-a.e. Moreover, the process $t \mapsto \tau'(t)$ has limits from the left, given by

$$\tau'(t-) := \int \mathbf{1}\{s < t\} D(e)\, \Phi'(d(s,e)), \quad t > 0.$$

Set $\tau'(0-) := 0$. Below we will write $\tau'(t) = \tau'(\Phi', t)$ and $\tau'(t-) = \tau'(\Phi', t-)$. Equation (6.7) also implies that $\nu(D > r) > 0$ for each $r > 0$, so that $\tau'(t) \to \infty$ as $t \to \infty$ holds $\mathbb{P}'$-a.s.

Motivated by [18, Proposition XII.(2.5)] we now define a process $W = (W_t)_{t \geq 0}$ as follows. Set $W_0 := 0$. Let $t > 0$. Then there exists $s > 0$ such that $\tau'(s-) \leq t < \tau'(s)$. By definition (6.8), there exists $e \in E$ such that $\Phi'\{(s, e)\} > 0$. Set

$$W_t := e(t - \tau'(s-)).$$

The process W is a measurable function of Φ'. We abuse the notation and write $W \equiv W(\Phi')$. By [18, Proposition XII.(2.5)] and the fact that (6.6) has the same distribution as Φ' it follows that $\mathbb{P}'_0(W \in \cdot) = \mathbb{P}_0(B^+ \in \cdot)$ is the distribution of a Brownian motion starting from 0. In fact, even more is true. Let $S := \inf\{t \geq 0 : B_t = 0\}$ and let B^S be the process B stopped at S; that is $B^S_t := B_t$ if $t \leq S$, and $B^S_t = 0$ otherwise. By the strong Markov property at S, the random measure

$$\sum_{s>0:\tau_{s-}<\tau_s} \delta_{(S+s, \epsilon_{\tau_{s-}})}$$

is for each $x \in \mathbb{R}$ a Poisson process on $(0, \infty) \times E$ under $\mathbb{P}'_x$ with intensity measure $dt\, \nu(de)$, independent of B^S. Define a process $B' = (B'_t)_{t \in \mathbb{R}}$ by

$$B'_t := \begin{cases} B_t, & \text{if } t \leq S, \\ W_{t-S}, & \text{if } t > S. \end{cases}$$

Then B' is a measurable function of (B^S, Φ') and we can write $B' \equiv B'(B^S, \Phi')$. Now we have

$$\mathbb{P}'_x(B' \in \cdot) = \mathbb{P}_x, \quad x \in \mathbb{R}. \tag{6.9}$$

Let $(s, e) \in (0, \infty) \times E$. A careful check of the definitions shows that

$$B'_t(B^S, \Phi' + \delta_{(s,e)}) = \begin{cases} B'_t, & \text{if } t \leq S + \tau'(s-), \\ e(t - S - \tau'(s-)), & \text{if } S + \tau'(s-) < t \leq S + \tau'(s-) + D(e), \\ B'_{t+D(e)}, & \text{if } S + \tau'(s-) + D(e) < t, \end{cases}$$

or

$$\theta_{S+\tau'(s-)} B'(B^S, \Phi' + \delta_{(s,e)}) = (\theta_{S+\tau'(s-)} B')^- \odot e \odot (\theta_{S+\tau'(s-)+D(e)} B')^+. \tag{6.10}$$

After these preparations, we can turn to the calculation of the Palm measure of N. Let $x \in \mathbb{R}$ and let $f \colon \Omega \to [0, \infty)$ be measurable. Then

$$\begin{aligned} \mathbb{E}_x \sum_{s \in L} f(\theta_s B) \mathbf{1}\{s \in (0, 1]\} &= \mathbb{E}_x \sum_{s:\tau_{s-}<\tau_s} f(\theta_{\tau_{s-}} B) \mathbf{1}\{\tau_{s-} \in (0, 1]\} \\ &= \mathbb{E}'_x \sum_{s:\tau'(s-)<\tau'(s)} f(\theta_{S+\tau'(s-)} B') \mathbf{1}\{S + \tau'(s-) \in (0, 1]\} \\ &= \mathbb{E}'_x \int f(\theta_{S+\tau'(s-)} B') \mathbf{1}\{S + \tau'(s-) \in (0, 1]\}\, \Phi'(ds \times E), \end{aligned}$$

where we have used (6.9) to get the second identity. Now we use the independence of B^S and Φ' along with the Mecke equation (see e.g. [7, Theorem 4.1]) to obtain that the last expression equals

$$\mathbb{E}'_x \int_E \int_0^\infty f(\theta_{S+\tau'(\Phi'+\delta_{(s,e)},s-)} B'(B^S, \Phi' + \delta_{(s,e)}) \mathbf{1}\{S + \tau'(\Phi' + \delta_{(s,e)}, s-) \in (0,1]\}\, ds\, \nu(de).$$

Clearly, we have $\tau'(\Phi' + \delta_{(s,e)}, s-) = \tau'(s-)$. So by (6.10), the above equals

$$\mathbb{E}'_x \int_E \int_0^\infty f((\theta_{S+\tau'(s)} B')^- \odot e \odot (\theta_{S+\tau'(s)} B')^+) \mathbf{1}\{S + \tau'(s) \in (0,1]\}\, ds\, \nu(de),$$

where we have used that $\int \mathbf{1}\{\tau'(s-) \neq \tau'(s)\}\, ds = 0$. Using (6.9) again gives

$$\begin{aligned}\mathbb{E}_x \sum_{s\in L} f(\theta_s B) \mathbf{1}\{s \in (0,1]\} &= \mathbb{E}_x \int_E \int_0^\infty f((\theta_{\tau_s} B)^- \odot e \odot (\theta_{\tau_s} B)^+) \mathbf{1}\{\tau_s \in (0,1]\}\, ds\, \nu(de) \\ &= \mathbb{E}_x \int_0^\infty \int_E f((\theta_t B)^- \odot e \odot (\theta_t B)^+) \mathbf{1}\{t \in (0,1]\}\, \nu(de)\, \ell^0(dt),\end{aligned}$$

where the second identity comes from a change of variables. Integrating with respect to x and using (6.5) (for $x = 0$) gives

$$\mathbb{E} \sum_{s\in L} f(\theta_s B) \mathbf{1}\{s \in (0,1]\} = \mathbb{E}_0 \int f(B^- \odot e \odot B^+)\, \nu(de),$$

which is the assertion. □

Proof of Theorem 6.3. Since ℓ^0 is diffuse and N_A is purely discrete, these two random measures are mutually singular. Moreover, (6.5) and Proposition 6.1 show that both random measures have intensity 1. Proposition 4.1, (6.5), and Corollary 6.2 imply the assertion. □

For $t \in \mathbb{R}$, let

$$G_t := \sup\{s \le t : B_s = 0\},$$

where $\sup \emptyset := -\infty$. Below we will write $G_t \equiv G_t(B)$. Also define $D_t := D_t(B) = \inf\{s > t : B_s = 0\}$. Note that $\mathbb{P}(G_t = D_t) = 0$ for each $t \in \mathbb{R}$. In particular,

$$\mathbb{P} = \mathbb{E}\mathbf{1}\{G_0 < D_0\} \mathbf{1}\{B \in \cdot\}. \tag{6.11}$$

By [16, Theorem (v)],

$$\mathbb{P} = \mathbb{E}_N \int_0^{D_0} \mathbf{1}\{\theta_t B \in \cdot\}\, dt,$$

implying that

$$\mathbb{P}(\theta_{G_0} \in \cdot) = \mathbb{E}_N \mathbf{1}\{B \in \cdot\} D_0; \tag{6.12}$$

see assertion (iv) of the above cited theorem. The measure

$$\nu' := \mathbb{P}(\epsilon_{G_0} \in \cdot) = \mathbb{P}(G_0 < D_0, \epsilon_{G_0} \in \cdot)$$

is known as *Bismut's excursion law*. It follows from (6.12) that $\nu'(de) = D(e)\,\nu(de)$.

Let $A \in \mathcal{E}$ satisfy $0 < \nu'(A) < \infty$. Similar to (6.1), define an invariant random measure N'_A by

$$N'_A := \nu'(A)^{-1} \int \mathbf{1}\{t \in \cdot\}\, \mathbf{1}\{\epsilon_{G_t} \in A\}\, dt. \tag{6.13}$$

It is easy to see that

$$\mathbb{P}_{N'_A}(B \in \cdot) = \nu'(A)^{-1}\mathbb{P}(B \in \cdot, \epsilon_{G_0} \in A). \tag{6.14}$$

We then have the following Bismut counterpart of Theorem 6.3.

Theorem 6.4. *Let $A \in \mathcal{E}$ be such that $0 < \nu'(A) < \infty$ and define the random measure N'_A by* (6.13). *Let*

$$T := \inf\{t > 0 : \ell^0[0,t] = N'_A[0,t]\}.$$

Then $\mathbb{P}_0(T < \infty) = 1$ and $\mathbb{P}_0(\theta_{G_T} B \in \cdot) = \mathbb{P}^- \odot \nu'(\cdot|A) \odot \mathbb{P}^+$.

Proof. Since ℓ^0 and Lebesgue measure are mutually singular, we can apply Theorem 3.1 and Proposition 4.1. This gives

$$\mathbb{P}_0(\theta_T B \in \cdot) = \mathbb{P}_{N'_A}(B \in \cdot) = \nu'(A)^{-1}\mathbb{P}(B \in \cdot, \epsilon_{G_0} \in A)$$

Since $G_t = G_0(\theta_t B) + t$, $t \in \mathbb{R}$, we hence have for each measurable $f\colon \Omega \to [0,\infty)$,

$$\begin{aligned}\mathbb{E}_0 f(\theta_{G_T} B) &= \mathbb{E}_0 f(\theta_{G_0(\theta_T B)}\theta_T B) = \mathbb{E}_{N'_A} f(\theta_{G_0} B)\\ &= \nu'(A)^{-1}\mathbb{E} f(\theta_{G_0} B)\mathbf{1}\{\epsilon_0(\theta_{G_0} B) \in A\},\end{aligned}$$

where we have used (6.14) and the identity $\epsilon_{G_0} = \epsilon_0(\theta_{G_0} B)$. Using (6.12) gives

$$\mathbb{E}_0 f(\theta_{G_T} B) = \nu'(A)^{-1}\mathbb{E}_N f(B) D_0 \mathbf{1}\{\epsilon_0 \in A\}.$$

By Proposition 6.1,

$$\mathbb{E}_0 f(\theta_{G_T} B) = \nu'(A)^{-1} \iiint f(w_1 \odot w_2 \odot w_3) D(w_2) \mathbf{1}\{w_2 \in A\}\, \mathbb{P}^-(dw_1)\, \nu(dw_2)\, \mathbb{P}^+(dw_3).$$

Using that $\nu'(de) = D(e)\,\nu(de)$, we obtain the assertion. □

Acknowledgments: We would like to thank Jim Pitman for some illuminating discussions of the topics in this paper. The first author thanks Steve Evans for supporting his visit to Berkeley and for giving valuable advice on some aspects of this work.

References

[1] D.J. Aldous and H. Thorisson (1993). Shift-coupling. *Stochastic Process. Appl.* **44**, 1-14.

[2] D. Geman and J. Horowitz (1973). Occupation times for smooth stationary processes. *Ann. Probab.* **1**, 131–137.

[3] A.E. Holroyd and T.M. Liggett (2001). How to find an extra head: optimal random shifts of Bernoulli and Poisson random fields. *Ann. Probab.* **29**, 1405-1425.

[4] A.E. Holroyd and Y. Peres (2005). Extra heads and invariant allocations. *Ann. Probab.* **33**, 31-52.

[5] K. Itô (1971). Poisson point processes attached to Markov processes. *Proc. 6th Berk. Symp. Math. Stat. Prob* **3**, 225-240.

[6] O. Kallenberg (2002). *Foundations of Modern Probability.* Second Edition, Springer, New York.

[7] G. Last and M.D. Penrose (2016). *Lectures on the Poisson Process.* Cambridge University Press, to appear. `http://www.math.kit.edu/stoch/~last/seite/lehrbuch_poissonp/de`

[8] G. Last, P. Mörters and H. Thorisson (2014). Unbiased shifts of Brownian motion. *Ann. Probab.* **42**, 431-463.

[9] G. Last and H. Thorisson (2009). Invariant transports of stationary random measures and mass-stationarity. *Ann. Probab.* **37**, 790-813.

[10] G. Last and H. Thorisson (2011). What is typical? *J. Appl. Probab.* **48A**, 379-390.

[11] T.M. Liggett (2002). Tagged particle distributions or how to choose a head at random. In *In and Out of Equlibrium* (V. Sidoravicious, ed.) 133-162, Birkhäuser, Boston.

[12] J. Mecke (1975). Invarianzeigenschaften allgemeiner Palmscher Maße. *Math. Nachr.* **65**, 335-344.

[13] P. Mörters and Y. Peres (2010). *Brownian Motion.* Cambridge University Press, Cambridge.

[14] P. Mörters, I. Redl (2016). Optimal embeddings by unbiased shifts of Brownian motion. arXiv:1605.07529.

[15] E. Perkins (1981). A global intrinsic characterisation of local time. *Ann. Probab.* **9**, 800–817.

[16] J. Pitman (1987). Stationary excursions. *Séminaire de Probabilités, XXI*, *Lecture Notes in Math.*, **1247**, 289-302, Springer, Berlin.

[17] J. Pitman and W. Tang (2015). The Slepian zero set, and Brownian bridge embedded in Brownian motion by a spacetime shift. *Electronic Journal of Probability*, **61**, 1-28.

[18] D. Revuz and M. Yor (1999). *Continuous Martingales and Brownian Motion.* Grundlehren der Mathematischen Wissenschaften **293**, Springer, Berlin.

[19] H. Thorisson (1996). Transforming random elements and shifting random fields. *Ann. Probab.* **24**, 2057-2064.

[20] H. Thorisson (2000). *Coupling, Stationarity, and Regeneration.* Springer, New York.